

# Acyclic Digraphs and Eigenvalues of $(0, 1)$ -Matrices

Brendan D. McKay, Department of Computer Science, Australian National University,  
Canberra, ACT 0200, AUSTRALIA

Frédérique E. Oggier<sup>1</sup>, Département de Mathématiques,  
Ecole Polytechnique Fédérale de Lausanne, 1015 Lausanne, SWITZERLAND

Gordon F. Royle, Department of Computer Science & Software Engineering, University of  
Western Australia, 35 Stirling Highway, Crawley, WA 6009, AUSTRALIA

N. J. A. Sloane<sup>2</sup>, Internet and Network Systems Research Department, AT&T Shannon Labs,  
180 Park Avenue, Florham Park, NJ 07932-0971, USA

Ian M. Wanless, Department of Computer Science, Australian National University,  
Canberra, ACT 0200, AUSTRALIA

Herbert S. Wilf, Mathematics Department, University of Pennsylvania,  
Philadelphia, PA 19104-6395, USA

October 24, 2003

## Abstract

We show that the number of acyclic directed graphs with  $n$  labeled vertices is equal to the number of  $n \times n$   $(0, 1)$ -matrices whose eigenvalues are positive real numbers.

Keywords:  $(0, 1)$ -matrix, acyclic, digraph, eigenvalue

AMS 2000 Classification: Primary 05A15, secondary 15A18, 15A36.

## 1. Weisstein's conjecture

A calculation was recently made by Eric W. Weisstein of Wolfram Research, Inc., to count the real  $n \times n$  matrices of 0's and 1's all of whose eigenvalues are real and positive. The resulting sequence of values, viz.,

$$1, 3, 25, 543, 29281$$

(for  $n = 1, 2, \dots, 5$ ) was then observed to coincide with the beginning of sequence [A003024](#) in [7], which counts acyclic digraphs with  $n$  labeled vertices. Weisstein conjectured that the sequences were in fact identical, and we prove this here.

Notation. A “digraph” means a graph with at most one edge directed from vertex  $i$  to vertex  $j$ , for  $1 \leq i \leq n, 1 \leq j \leq n$ . Loops and cycles of length two are permitted, but parallel edges are forbidden. “Acyclic” means there are no cycles of any length.

**Theorem 1.** *For each  $n = 1, 2, 3, \dots$ , the number of acyclic directed graphs with  $n$  labeled vertices is equal to the number of  $n \times n$  matrices of 0's and 1's whose eigenvalues are positive real numbers.*

<sup>1</sup>This work was carried out during F. E. Oggier's visit to AT&T Shannon Labs during the summer of 2003. She thanks the Fonds National Suisse, Bourses et Programmes d'Échange for support.

<sup>2</sup>To whom correspondence should be addressed. [Email: njas@research.att.com, phone: 973 360 8415, fax: 973 360 8178.]

**Proof.** Suppose we are given an acyclic directed graph  $G$ . Let  $A = A(G)$  be its vertex adjacency matrix. Then  $A$  has only 0's on the diagonal, else cycles of length 1 would be present. So define  $B = I + A$ , and note that  $B$  is also a matrix of 0's and 1's. We claim  $B$  has only positive eigenvalues.

Indeed, the eigenvalues will not change if we renumber the vertices of the graph  $G$  consistently with the partial order that it generates. But then  $A = A(G)$  would be strictly upper triangular, and  $B$  would be upper triangular with 1's on the diagonal. Hence all of its eigenvalues are equal to 1.

Conversely, let  $B$  be a  $(0, 1)$ -matrix whose eigenvalues are all positive real numbers. Then we have

$$\begin{aligned}
1 &\geq \frac{1}{n} \text{Trace}(B) && (\text{since all } B_{i,i} \leq 1) \\
&= \frac{1}{n} (\lambda_1 + \lambda_2 + \dots + \lambda_n) \\
&\geq (\lambda_1 \lambda_2 \dots \lambda_n)^{\frac{1}{n}} && (\text{by the arithmetic-geometric mean inequality}) \\
&= (\det B)^{\frac{1}{n}} \\
&\geq 1 && (\text{since } \det B \text{ is a positive integer}).
\end{aligned} \tag{1}$$

Since the arithmetic and geometric means of the eigenvalues are equal, the eigenvalues are all equal, and in fact all  $\lambda_i(B) = 1$ .

Now regard  $B$  as the adjacency matrix of a digraph  $H$ , which has a loop at each vertex. Since

$$\text{Trace}(B^k) = \sum_{i=1}^n \lambda_i^k = \sum_{i=1}^n 1 = n,$$

for all  $k$ , the number of closed walks in  $H$ , of each length  $k$ , is  $n$ .

Since the trace of  $B$  is equal to  $n$ , all diagonal entries of  $B$  are 1's. Thus we account for all  $n$  of the closed walks of length  $k$  that exist in the graph  $H$  by the loops at each vertex. There are no closed walks of any length that use an edge of  $H$  other than the loops at the vertices.

Put  $A = B - I$ . Then  $A$  is a  $(0, 1)$ -matrix that is the adjacency matrix of an acyclic digraph.  $\square$

Remark. The only related result we have found in the literature is the theorem [3, p. 81] that a digraph  $G$  contains no cycle if and only if all eigenvalues of the adjacency matrix are 0.

## 2. Corollaries.

The proof also establishes the following results.

(i) Let  $B$  be a  $(0, 1)$ -matrix whose eigenvalues are all positive real numbers. Then the eigenvalues are in fact all equal to 1. The only symmetric  $(0, 1)$ -matrix with positive eigenvalues is the identity.

(ii) Let  $B$  be an  $n \times n$  matrix with integer entries and  $\text{Trace}(B) \leq n$ . Then  $B$  has all eigenvalues real and positive if and only if  $B = I + N$ , where  $N$  is nilpotent.

(iii) If a digraph contains a cycle, then its adjacency matrix has an eigenvalue which is zero, negative, or strictly complex. In fact, a more detailed argument, not given here, shows that if the length of the shortest cycle is at least 3, then there is a strictly complex eigenvalue.

(iv) The eigenvalues of a digraph consist of  $n - k$  0's and  $k$  1's if and only if the digraph is acyclic apart from  $k$  loops.

(v) Define two matrices  $B_1, B_2$  to be *equivalent* if there is a permutation matrix  $P$  such that  $P'B_1P = B_2$ . Then the number of equivalence classes of  $n \times n$   $(0, 1)$ -matrices with all eigenvalues

positive is equal to the number of acyclic digraphs with  $n$  unlabeled vertices. (These numbers form sequence [A003087](#) in [7].)

Proof. Two labeled graphs  $G_1, G_2$  with adjacency matrices  $A(G_1), A(G_2)$  correspond to the same unlabeled graph if and only if there is a permutation matrix  $P$  such that  $P'A(G_1)P = A(G_2)$ . The result now follows immediately from the theorem.  $\square$

(vi) Let  $B$  be an  $n \times n$   $(-1, +1)$ -matrix with all eigenvalues real and positive. Then  $n = 1$  and  $B = [1]$ .

Proof. The argument that led to (1) still applies and shows that all the eigenvalues are 1,  $\det B = 1$  and  $\text{Trace}(B) = n$ . By adding or subtracting the first row of  $B$  from all other rows we can clear the first column, obtaining a matrix

$$C = \begin{bmatrix} 1 & * \\ \mathbf{0} & D \end{bmatrix},$$

where  $\mathbf{0}$  is a column of 0's and  $D$  is an  $(n-1) \times (n-1)$  matrix with entries  $-2, 0, +2$  and  $\det D = \det C = \det B = 1$ . Hence  $2^{n-1}$  divides 1, so  $n = 1$ .  $\square$

It would be interesting to investigate the connections between matrices and graphs in other cases—for example if the eigenvalues are required only to be real and nonnegative (see sequences [A086510](#), [A087488](#) in [7] for the initial values), or if the entries are  $-1, 0$  or  $1$  ([A085506](#)).

### 3. Bibliographic remarks

Acyclic digraphs were first counted by Robinson [5, 6], and independently by Stanley [8]: if  $R_n$  is the number of acyclic digraphs with  $n$  labeled vertices, then

$$R_n = \sum_{k=1}^n (-1)^{k+1} \binom{n}{k} 2^{k(n-k)} R_{n-k},$$

for  $n \geq 1$ , with  $R_0 = 1$ , and

$$\sum_{n=0}^{\infty} R_n \frac{x^n}{2^{\binom{n}{2}} n!} = \left[ \sum_{n=0}^{\infty} (-1)^n \frac{x^n}{2^{\binom{n}{2}} n!} \right]^{-1}.$$

The asymptotic behavior is

$$R_n \sim n! \frac{2^{\binom{n}{2}}}{M p^n},$$

where  $p = 1.488\dots$  and  $M = 0.474\dots$

The asymptotic behavior of  $R(n, q)$ , the number of these graphs that have  $q$  edges, was found by Bender *et al.* [1, 2], and the number that have specified numbers of sources and sinks has been found by Gessel [4].

### References

- [1] E. A. Bender, L. B. Richmond, R. W. Robinson and N. C. Wormald, The asymptotic number of acyclic digraphs, I, *Combinatorica* **6** (1986), 15–22.
- [2] E. A. Bender and R. W. Robinson, The asymptotic number of acyclic digraphs, II, *J. Combin. Theory*, Ser. **B** **44** (1988), 363–369.

- [3] D. M. Cvetković, M. Doob and H. Sachs, *Spectra of Graphs*, third ed., Barth, Heidelberg, 1995.
- [4] I. M. Gessel, Counting acyclic digraphs by sources and sinks, *Discrete Math.*, **160** (1996), 253–258.
- [5] R. W. Robinson, Enumeration of acyclic digraphs, in: R. C. Bose *et al.* (Eds.), *Proc. Second Chapel Hill Conf. on Combinatorial Mathematics and its Applications (Univ. North Carolina, Chapel Hill, N.C., 1970)*, Univ. North Carolina, Chapel Hill, N.C., 1970, pp. 391–399.
- [6] R. W. Robinson, Counting labeled acyclic digraphs, in: F. Harary (Ed.), *New Directions in the Theory of Graphs*, Academic Press, NY, 1973, pp. 239–273.
- [7] N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences*, published electronically at [www.research.att.com/~njas/sequences/](http://www.research.att.com/~njas/sequences/), 1996–2003.
- [8] R. P. Stanley, Acyclic orientations of graphs, *Discrete Math.*, **5** (1973), 171–178.